

NOR-TEST

Norwegian Tenecteplase Stroke Trial

A randomized controlled trial of
tenecteplase vs. alteplase
in acute ischemic stroke

Nicola Logallo MD PhD

Hypothesis & Methods

NOR-TEST

Norwegian Tenecteplase Stroke Trial

Hypothesis

- TNK 0,4 mg/kg has superior efficacy
- TNK 0,4 mg/kg is safe

Design

- Randomized Open label Blinded Endpoints
- Tenecteplase 0.4 mg/kg vs. alteplase 0.9 mg/kg
- Randomisation 1:1

Inclusion criteria

- Ischemic stroke patients > 18 years
- Mesurable NIHSS deficit
- Treatment start <4.5 hours

Study endpoints

NOR-TEST

Norwegian Tenecteplase Stroke Trial

- Primary endpoint
 - Outcome at 3 months (mRS 0-1)
- Secondary endpoints
 - Safety (SICH / ICH)
 - NIHSS at 24 hours
 - mRS 3 months (ordinal shift)

NOR-TEST

Norwegian Tenecteplase Stroke Trial



NORSTROKE

Norwegian Stroke Research Co-operation

Bodö

Trondheim

Molde

Förde

Bergen

Haugesund

Stavanger

Tönsberg

Skien

Drammen

Oslo

Akershus

Baerum